

Package ‘VBMS’

October 8, 2025

Title Variational Bayesian Algorithm for Multi-Source Heterogeneous Models

Version 1.0.0

Description A Variational Bayesian algorithm for high-dimensional multi-source heterogeneous linear models. More details have been written up in a paper submitted to the journal Statistics in Medicine, and the details of variational Bayesian methods can be found in Ray and Szabo (2021) <[doi:10.1080/01621459.2020.1847121](https://doi.org/10.1080/01621459.2020.1847121)>. It simultaneously performs parameter estimation and variable selection. The algorithm supports two model settings: (1) local models, where variable selection is only applied to homogeneous coefficients, and (2) global models, where variable selection is also performed on heterogeneous coefficients. Two forms of Spike-and-Slab priors are available: the Laplace distribution and the Gaussian distribution as the Slab component.

License MIT + file LICENSE

Imports pracma, selectiveInference, MASS

Encoding UTF-8

RoxygenNote 7.3.2

NeedsCompilation no

Author Lu Luo [aut, cre],
Huiqiong Li [aut]

Maintainer Lu Luo <luolu@stu.ynu.edu.cn>

Repository CRAN

Date/Publication 2025-10-08 19:40:13 UTC

Contents

vbms	2
vb_gauss_global	3
vb_gauss_local	4
vb_lap_global	5
vb_lap_local	6

vbms*Variational Bayesian Algorithm for Multi-Source Heterogeneous Models.*

Description

This package implements a variational Bayesian algorithm for high-dimensional multi-source heterogeneous linear models. It simultaneously performs parameter estimation and variable selection. The algorithm supports two model settings: (1) local models, where variable selection is only applied to homogeneous coefficients, and (2) global models, where variable selection is also performed on heterogeneous coefficients. Two forms of Spike-and-Slab priors are available: the Laplace distribution and the Gaussian distribution as the Slab component.

Usage

```
vbms(
  X,
  Z,
  Y,
  global,
  prior,
  max_iter = 1000,
  tol = 1e-06,
  a = 1,
  b = 10,
  lambda = 1
)
```

Arguments

X	Homogeneous covariates
Z	Heterogeneous covariates
Y	Response covariates
global	Indicates whether variable selection is required for het coefficients, if TRUE, Variable selection will be made for het coefficients.
prior	Forms of Slab distribution in Spike-and-Slab prior, "laplace" or "gauss".
max_iter	Maximum number of iterations, Default:1000
tol	Algorithm convergence tolerance, Default:1e-6
a	A prior of Beta distribution, Default:1
b	A prior of Beta distribution, Default:10
lambda	A prior of Laplace distribution, Default:1

Value

mu_hom	The mean of the homogeneous coefficients
sigma_hom	The variance of homogeneous coefficients
gamma_hom	Selection indicators for homogeneous coefficients
mu_het	The mean of the heterogeneous coefficients
sigma_het	The variance of heterogeneous coefficients
gamma_het	Selection indicators for heterogeneous coefficients (NULL for local models)

Examples

```

# Simulate multi-source heterogeneous data
n <- 50 # number of samples per source
K <- 3 # number of sources
p <- 100 # number of homogeneous covariates
q <- 5 # number of heterogeneous covariates

set.seed(1)
theta <- matrix(c(c(-1,0.5,1,-0.5,2),rep(0,p-5)), ncol = 1)
beta <- matrix(1, nrow = q, ncol = K)
for (k in 1:K) {
  beta[,k] <- matrix(c(rep(log(k+1),5),rep(0,q-5)), ncol = 1)
}

zdata <- MASS::mvrnorm(K*n, rep(0,q), diag(q))
Z <- array(data=zdata,dim=c(n,q,K))
xdata <- MASS::mvrnorm(K*n, rep(0,p), diag(p))
X <- array(data=xdata,dim=c(n,p,K))
Y <- matrix(0, nrow = n, ncol = K)

for (k in 1:K) {
  Y[,k] <- MASS::mvrnorm(1, X[, ,k] %*% theta + Z[, ,k] %*% beta[,k], diag(n))
}

# Fit local model with Laplace prior
res <- vbms(X, Z, Y, global=FALSE, prior='laplace')

# View results
print(head(res$mu_hom))      # Homogeneous coefficients mean
print(head(res$gamma_hom))    # Homogeneous variable selection
print(res$mu_het)            # Heterogeneous coefficients mean

```

Description

A variational Bayesian algorithm is proposed for multi-source heterogeneous models under the Laplace Spike-and-Slab prior, enabling simultaneous variable selection for both homogeneous and #' heterogeneous covariates.

Usage

```
vb_gauss_global(
  X,
  Z,
  Y,
  max_iter = 1000,
  tol = 1e-06,
  a = 1,
  b = 10,
  lambda = 1
)
```

Arguments

X	Homogeneous covariates
Z	Heterogeneous covariates
Y	Response covariates
max_iter	Maximum number of iterations, Default:1000
tol	Algorithm convergence tolerance, Default:1e-6
a	A prior of Beta distribution, Default:1
b	A prior of Beta distribution, Default:10
lambda	A prior of Laplace distribution, Default:1

Value

The mean of the homogeneity coefficient:mu1; The variance of homogeneity coefficient:sigma1; Selection coefficient:gamma1; The mean of the heterogeneous coefficient:mu2; The variance of heterogeneous coefficient:sigma2; Selection heterogeneous:gamma2.

Description

A variational Bayesian algorithm, based on the Gauss Spike-and-Slab prior, is tailored for multi-source heterogeneous models and focuses on variable selection exclusively for the homogeneous covariates.

Usage

```
vb_gauss_local(X, Z, Y, max_iter, tol, a = 1, b = 10, lambda = 1)
```

Arguments

X	Homogeneous covariates
Z	Heterogeneous covariates
Y	Response covariates
max_iter	Maximum number of iterations, Default:1000
tol	Algorithm convergence tolerance, Default:1e-6
a	A prior of Beta distribution, Default:1
b	A prior of Beta distribution, Default:10
lambda	A prior of Laplace distribution, Default:1

Value

The mean of the homogeneity coefficient:mu; The variance of homogeneity coefficient:sigma; Selection coefficient:gamma; Mean and covariance of heterogeneity coefficients:m, s2.

vb_lap_global

*Global Laplace VB***Description**

A variational Bayesian algorithm is proposed for multi-source heterogeneous models under the Laplace Spike-and-Slab prior, enabling simultaneous variable selection for both homogeneous and #' heterogeneous covariates.

Usage

```
vb_lap_global(X, Z, Y, max_iter = 1000, tol = 1e-06, a = 1, b = 10, lambda = 1)
```

Arguments

X	Homogeneous covariates
Z	Heterogeneous covariates
Y	Response covariates
max_iter	Maximum number of iterations, Default:1000
tol	Algorithm convergence tolerance, Default:1e-6
a	A prior of Beta distribution, Default:1
b	A prior of Beta distribution, Default:10
lambda	A prior of Laplace distribution, Default:1

Value

The mean of the homogeneity coefficient:mu1; The variance of homogeneity coefficient:sigma1; Selection coefficient:gamma1; The mean of the heterogeneous coefficient:mu2; The variance of heterogeneous coefficient:sigma2; Selection heterogeneous:gamma2.

vb_lap_local

*Local Laplace VB***Description**

A variational Bayesian algorithm, based on the Laplace Spike-and-Slab prior, is tailored for multi-source heterogeneous models and focuses on variable selection exclusively for the homogeneous covariates.

Usage

```
vb_lap_local(X, Z, Y, max_iter = 1000, tol = 1e-06, a = 1, b = 10, lambda = 1)
```

Arguments

X	Homogeneous covariates
Z	Heterogeneous covariates
Y	Response covariates
max_iter	Maximum number of iterations, Default:1000
tol	Algorithm convergence tolerance, Default:1e-6
a	A prior of Beta distribution, Default:1
b	A prior of Beta distribution, Default:10
lambda	A prior of Laplace distribution, Default:1

Value

The mean of the homogeneity coefficient:mu; The variance of homogeneity coefficient:sigma; Selection coefficient:gamma; Mean and covariance of heterogeneity coefficients:m, s2.

Index

`vb_gauss_global`, 3
`vb_gauss_local`, 4
`vb_lap_global`, 5
`vb_lap_local`, 6
`vbms`, 2